

## CHEN INEQUALITIES FOR SUBMANIFOLDS OF A COSYMPLECTIC SPACE FORM WITH A SEMI-SYMMETRIC METRIC CONNECTION

BY

CİHAN ÖZGÜR and CENGİZHAN MURATHAN

**Abstract.** In this paper, we prove Chen inequalities for submanifolds of a cosymplectic space form of constant  $\varphi$ -sectional curvature  $N^{2m+1}(c)$  endowed with a semi-symmetric metric connection, i.e., relations between the mean curvature associated with the semi-symmetric metric connection, scalar and sectional curvatures,  $k$ -Ricci curvature and the sectional curvature of the ambient space.

**Mathematics Subject Classification 2000:** 53C40, 53B05, 53B15.

**Key words:** semi-symmetric metric connection, Chen inequality, cosymplectic space form, Ricci curvature.

### 1. Introduction

The idea of a semi-symmetric linear connection on a differentiable manifold was introduced by FRIEDMANN and SCHOUTENN in [10]. The notion of a semi-symmetric metric connection on a Riemannian manifold was introduced by HAYDEN in [11]. Later in [22], YANO studied some properties of a Riemannian manifold endowed with a semi-symmetric metric connection. In the case of hypersurfaces, in [12] and [13], IMAI found some properties of a Riemannian manifold and a hypersurface of a Riemannian manifold with a semi-symmetric metric connection, respectively. In [20], NAKAO studied submanifolds of a Riemannian manifold with semi-symmetric connections.

In [5], CHEN recalled that one of the basic interests of submanifold theory is to establish simple relationships between the main extrinsic invariants and the main intrinsic invariants of a submanifold. Many famous results

in differential geometry can be regarded as results in this respect. The main extrinsic invariant is the squared mean curvature and the main intrinsic invariants include the classical curvature invariants namely the scalar curvature, the sectional curvature or the Ricci curvature. There are also other important modern intrinsic invariants of submanifolds introduced by CHEN [8].

Afterwards, many geometers studied similar problems for different submanifolds in various ambient spaces, for example see [3], [4], [7], [9], [16], [17] and [21].

In [14] and [23], submanifolds of cosymplectic space forms satisfying Chen's inequalities were studied.

Recently, in [18] and [19], the first author and MIHAI proved Chen inequalities for submanifolds of real space forms with a semi-symmetric metric connection and Chen inequalities for submanifolds of complex space forms and Sasakian space forms endowed with semi-symmetric metric connections, respectively.

Motivated by the studies of the above authors, in this study, we consider Chen inequalities for submanifolds in cosymplectic space forms of constant  $\varphi$ -sectional curvature  $N^{2m+1}(c)$  endowed with a semi-symmetric metric connection.

## 2. Semi-symmetric metric connection

Let  $N^{n+p}$  be an  $(n+p)$ -dimensional Riemannian manifold and  $\tilde{\nabla}$  a linear connection on  $N^{n+p}$ . If the torsion tensor  $\tilde{T}$  of  $\tilde{\nabla}$ , defined by

$$\tilde{T}(\tilde{X}, \tilde{Y}) = \tilde{\nabla}_{\tilde{X}}\tilde{Y} - \tilde{\nabla}_{\tilde{Y}}\tilde{X} - [\tilde{X}, \tilde{Y}],$$

for any vector fields  $\tilde{X}$  and  $\tilde{Y}$  on  $N^{n+p}$ , satisfies

$$\tilde{T}(\tilde{X}, \tilde{Y}) = \omega(\tilde{Y})\tilde{X} - \omega(\tilde{X})\tilde{Y}$$

for a 1-form  $\omega$ , then the connection  $\tilde{\nabla}$  is called a *semi-symmetric connection*.

Let  $g$  be a Riemannian metric on  $N^{n+p}$ . If  $\tilde{\nabla}g = 0$ , then  $\tilde{\nabla}$  is called a *semi-symmetric metric connection* on  $N^{n+p}$ .

A semi-symmetric metric connection  $\tilde{\nabla}$  on  $N^{n+p}$  is given by

$$\tilde{\nabla}_{\tilde{X}}\tilde{Y} = \overset{\circ}{\nabla}_{\tilde{X}}\tilde{Y} + \omega(\tilde{Y})\tilde{X} - g(\tilde{X}, \tilde{Y})U,$$

for any vector fields  $\tilde{X}$  and  $\tilde{Y}$  on  $N^{n+p}$ , where  $\overset{\circ}{\nabla}$  denotes the Levi-Civita connection with respect to the Riemannian metric  $g$  and  $U$  is a vector field defined by  $g(U, \tilde{X}) = \omega(\tilde{X})$ , for any vector field  $\tilde{X}$  [22].

We will consider a Riemannian manifold  $N^{n+p}$  endowed with a semi-symmetric metric connection  $\tilde{\nabla}$  and the Levi-Civita connection denoted by  $\overset{\circ}{\nabla}$ .

Let  $M^n$  be an  $n$ -dimensional submanifold of an  $(n+p)$ -dimensional Riemannian manifold  $N^{n+p}$ . On the submanifold  $M^n$  we consider the induced semi-symmetric metric connection denoted by  $\nabla$  and the induced Levi-Civita connection denoted by  $\overset{\circ}{\nabla}$ .

Let  $\tilde{R}$  be the curvature tensor of  $N^{n+p}$  with respect to  $\tilde{\nabla}$  and  $\overset{\circ}{R}$  the curvature tensor of  $N^{n+p}$  with respect to  $\overset{\circ}{\nabla}$ . We also denote by  $R$  and  $\overset{\circ}{R}$  the curvature tensors of  $\nabla$  and  $\overset{\circ}{\nabla}$ , respectively, on  $M^n$ .

The Gauss formulas with respect to  $\nabla$ , respectively  $\overset{\circ}{\nabla}$  can be written as:

$$\begin{aligned}\tilde{\nabla}_X Y &= \nabla_X Y + h(X, Y), \quad X, Y \in \chi(M), \\ \overset{\circ}{\nabla}_X Y &= \overset{\circ}{\nabla}_X Y + \overset{\circ}{h}(X, Y), \quad X, Y \in \chi(M),\end{aligned}$$

where  $\overset{\circ}{h}$  is the second fundamental form of  $M^n$  in  $N^{n+p}$  and  $h$  is a  $(0, 2)$ -tensor on  $M^n$ . According to the formula (7) from [20]  $h$  is also symmetric. The Gauss equation for the submanifold  $M^n$  into an  $(n+p)$ -dimensional Riemannian manifold  $N^{n+p}$  is

$$\begin{aligned}(2.1) \quad \tilde{R}(X, Y, Z, W) &= \overset{\circ}{R}(X, Y, Z, W) + g(\overset{\circ}{h}(X, Z), \overset{\circ}{h}(Y, W)) \\ &\quad - g(\overset{\circ}{h}(X, W), \overset{\circ}{h}(Y, Z)).\end{aligned}$$

One denotes by  $\overset{\circ}{H}$  the mean curvature vector of  $M^n$  in  $N^{n+p}$ .

Then the curvature tensor  $\tilde{R}$  with respect to the semi-symmetric metric connection  $\tilde{\nabla}$  on  $N^{n+p}$  can be written as (see [13])

$$\begin{aligned}(2.2) \quad \tilde{R}(X, Y, Z, W) &= \overset{\circ}{R}(X, Y, Z, W) - \alpha(Y, Z)g(X, W) \\ &\quad + \alpha(X, Z)g(Y, W) - \alpha(X, W)g(Y, Z) + \alpha(Y, W)g(X, Z),\end{aligned}$$

for any vector fields  $X, Y, Z, W \in \chi(M^n)$ , where  $\alpha$  is a  $(0, 2)$ -tensor field defined by

$$\alpha(X, Y) = (\overset{\circ}{\nabla}_X \omega)Y - \omega(X)\omega(Y) + \frac{1}{2}\omega(P)g(X, Y), \quad \forall X, Y \in \chi(M).$$

Denote by  $\lambda$  the trace of  $\alpha$ .

Let  $\pi \subset T_x M^n$ ,  $x \in M^n$ , be a 2-plane section. Denote by  $K(\pi)$  the sectional curvature of  $M^n$  with respect to the induced semi-symmetric metric connection  $\nabla$ . For any orthonormal basis  $\{e_1, \dots, e_m\}$  of the tangent space  $T_x M^n$ , the scalar curvature  $\tau$  at  $x$  is defined by

$$\tau(x) = \sum_{1 \leq i < j \leq n} K(e_i \wedge e_j).$$

Recall that the *Chen first invariant* is given by

$$\delta_M(x) = \tau(x) - \inf \{K(\pi) \mid \pi \subset T_x M^n, x \in M^n, \dim \pi = 2\},$$

(see, for example, [8]), where  $M^n$  is a Riemannian manifold,  $K(\pi)$  is the sectional curvature of  $M^n$  associated with a 2-plane section,  $\pi \subset T_x M^n$ ,  $x \in M^n$  and  $\tau$  is the scalar curvature at  $x$ .

The following algebraic Lemma is well-known.

**Lemma 2.1** ([5]). *Let  $a_1, a_2, \dots, a_n, b$  be  $(n + 1)$  ( $n \geq 2$ ) real numbers such that*

$$\left( \sum_{i=1}^n a_i \right)^2 = (n - 1) \left( \sum_{i=1}^n a_i^2 + b \right).$$

*Then  $2a_1 a_2 \geq b$ , with equality holding if and only if  $a_1 + a_2 = a_3 = \dots = a_n$ .*

Let  $M^n$  be an  $n$ -dimensional Riemannian manifold,  $L$  a  $k$ -plane section of  $T_x M^n$ ,  $x \in M^n$ , and  $X$  a unit vector in  $L$ .

We choose an orthonormal basis  $\{e_1, \dots, e_k\}$  of  $L$  such that  $e_1 = X$ .

One defines [7] the *Ricci curvature* (or *k-Ricci curvature*) of  $L$  at  $X$  by

$$Ric_L(X) = K_{12} + K_{13} + \dots + K_{1k},$$

where  $K_{ij}$  denotes, as usual, the sectional curvature of the 2-plane section spanned by  $e_i, e_j$ . For each integer  $k$ ,  $2 \leq k \leq n$ , the Riemannian invariant  $\Theta_k$  on  $M^n$  is defined by:

$$\Theta_k(x) = \frac{1}{k-1} \inf_{L, X} Ric_L(X), \quad x \in M^n,$$

where  $L$  runs over all  $k$ -plane sections in  $T_x M^n$  and  $X$  runs over all unit vectors in  $L$ .

### 3. Chen first inequality for submanifolds of cosymplectic manifolds

Let  $N^{2m+1}$  be a  $(2m+1)$ -dimensional almost contact manifold endowed with an almost contact structure  $(\varphi, \xi, \eta)$ , that is,  $\varphi$  is a  $(1,1)$ -tensor field,  $\xi$  is a vector field and  $\eta$  is 1-form such that  $\varphi^2 X = -X + \eta(X)\xi$ ,  $\eta(\xi) = 1$ . Then,  $\varphi\xi = 0$  and  $\eta \circ \varphi = 0$ . The almost contact structure is said to be *normal* if the induced almost complex structure  $J$  on the product manifold  $N \times \mathbb{R}$  defined by  $J(X, \lambda \frac{d}{dt}) = (\varphi X - \lambda\xi, \eta(X) \frac{d}{dt})$  is integrable, where  $X$  is tangent to  $N$ ,  $t$  the coordinate of  $\mathbb{R}$  and  $\lambda$  a smooth function on  $N \times \mathbb{R}$ . The condition for being normal is equivalent to vanishing of the torsion tensor  $[\varphi, \varphi] + 2d\eta \otimes \xi$ , where  $[\varphi, \varphi]$  is the Nijenhuis tensor of  $\varphi$ .

Let  $g$  be a compatible Riemannian metric with  $(\varphi, \xi, \eta)$ , that is,  $g(\varphi X, \varphi Y) = g(X, Y) - \eta(X)\eta(Y)$  or equivalently,  $\Phi(X, Y) = g(X, \varphi Y) = -g(\varphi X, Y)$  and  $g(X, \xi) = \eta(X)$  for all  $X, Y \in TN$ . Then  $N$  becomes an almost contact metric manifold equipped with an almost contact metric structure  $(\varphi, \xi, \eta, g)$ . If  $\Phi = d\eta$ , the almost contact structure is a contact structure. A normal contact structure such that the fundamental 2-form  $\Phi$  and 1-form  $\eta$  are closed is called a *cosymplectic structure*. It can be shown that the cosymplectic structure is characterized by

$$(3.1) \quad \overset{\circ}{\nabla}_X \varphi = 0 \quad \text{and} \quad \overset{\circ}{\nabla}_X \eta = 0,$$

(see [2]). From formula (3.1) it follows that  $\overset{\circ}{\nabla}_X \xi = 0$ .

A cosymplectic manifold  $N^{2m+1}$  is said to be a *cosymplectic space form* [15] if the  $\varphi$ -sectional curvature is constant  $c$  along  $N^{2m+1}$ . A cosymplectic space form will be denoted by  $N^{2m+1}(c)$ . Then the curvature tensor  $\tilde{R}$  of  $N^{2m+1}(c)$  can be expressed as

$$(3.2) \quad \begin{aligned} \overset{\circ}{\tilde{R}}(X, Y, Z, W) = & \frac{c}{4} [g(X, W)g(Y, Z) - g(X, Z)g(Y, W) \\ & + g(X, \varphi W)g(Y, \varphi Z) - g(X, \varphi Z)g(Y, \varphi W) - 2g(X, \varphi Y)g(Z, \varphi W) \\ & - \eta(Y)\eta(Z)g(X, W) + \eta(Y)\eta(W)g(X, Z) \\ & - \eta(X)\eta(W)g(Y, Z) + \eta(X)\eta(Z)g(Y, W)]. \end{aligned}$$

If  $N^{2m+1}(c)$  is a cosymplectic space form of constant  $\varphi$ -sectional curvature  $c$  with a semi-symmetric metric connection then from (2.2) and (3.2) it follows that

$$\begin{aligned}
 \tilde{R}(X, Y, Z, W) &= \frac{c}{4}[g(X, W)g(Y, Z) - g(X, Z)g(Y, W) \\
 &+ g(X, \varphi W)g(Y, \varphi Z) - g(X, \varphi Z)g(Y, \varphi W) - 2g(X, \varphi Y)g(Z, \varphi W) \\
 (3.3) \quad &- \eta(Y)\eta(Z)g(X, W) + \eta(Y)\eta(W)g(X, Z) \\
 &- \eta(X)\eta(W)g(Y, Z) + \eta(X)\eta(Z)g(Y, W)] \\
 &- \alpha(Y, Z)g(X, W) + \alpha(X, Z)g(Y, W) \\
 &- \alpha(X, W)g(Y, Z) + \alpha(Y, W)g(X, Z).
 \end{aligned}$$

Let  $M^n, n \geq 3$ , be an  $n$ -dimensional submanifold of an  $(2m + 1)$ -dimensional cosymplectic manifold  $N^{n+p}(c)$  of constant  $\varphi$ -sectional curvature  $c$ . For any tangent vector field  $X$  to  $M^n$ , we put

$$\varphi X = PX + FX,$$

where  $PX$  and  $FX$  are tangential and normal components of  $\varphi X$ , respectively and we decompose

$$\xi = \xi^\top + \xi^\perp,$$

where  $\xi^\top$  and  $\xi^\perp$  denotes the tangential and normal parts of  $\xi$ .

Denote by  $\Theta^2(\pi) = g^2(Pe_1, e_2)$ , where  $\{e_1, e_2\}$  is an orthonormal basis of a 2-plane section  $\pi$ , is a real number in  $[0, 1]$ , independent of the choice of  $e_1, e_2$  (see [1]).

For submanifolds of a cosymplectic space form  $N^{2m+1}(c)$  of constant  $\varphi$ -sectional curvature  $c$  endowed with a semi-symmetric metric connection we establish the following optimal inequality.

**Theorem 3.1.** *Let  $M^n, n \geq 3$ , be an  $n$ -dimensional submanifold of an  $(2m + 1)$ -dimensional cosymplectic space form  $N^{2m+1}(c)$  of constant  $\varphi$ -sectional curvature  $c$  endowed with a semi-symmetric metric connection  $\bar{\nabla}$ . We have:*

$$\begin{aligned}
 (3.4) \quad \tau(x) - K(\pi) &\leq (n - 2) \left[ \frac{n^2}{2(n - 1)} \|H\|^2 + (n + 1) \frac{c}{8} - \lambda \right] \\
 &- \frac{c}{4} (3\Theta^2(\pi) - \frac{3}{2} \|P\|^2 + (n - 1) \|\xi^\top\|^2 - \|\xi_\pi\|^2) - \text{trace}(\alpha|_{\pi^\perp}),
 \end{aligned}$$

where  $\pi$  is a 2-plane section of  $T_x M^n, x \in M^n$ .

**Proof.** From [20], the Gauss equation with respect to the semi-symmetric metric connection is

$$(3.5) \quad \begin{aligned} \tilde{R}(X, Y, Z, W) &= R(X, Y, Z, W) + g(h(X, Z), h(Y, W)) \\ &\quad - g(h(Y, Z), h(X, W)). \end{aligned}$$

Let  $x \in M^n$  and  $\{e_1, e_2, \dots, e_n\}$  and  $\{e_{n+1}, \dots, e_{2m+1}\}$  be orthonormal basis of  $T_x M^n$  and  $T_x^\perp M^n$ , respectively. For  $X = W = e_i, Y = Z = e_j, i \neq j$ , from the equation (3.3) it follows that:

$$(3.6) \quad \begin{aligned} \tilde{R}(e_i, e_j, e_j, e_i) &= \frac{c}{4} + \frac{3c}{4}g^2(Pe_j, e_i) - \frac{c}{4}\{\eta(e_i)^2 + \eta(e_j)^2\} \\ &\quad - \alpha(e_i, e_i) - \alpha(e_j, e_j). \end{aligned}$$

From (3.5) and (3.6) we get

$$\begin{aligned} &\frac{c}{4} + \frac{3c}{4}g^2(Pe_j, e_i) - \frac{c}{4}\{\eta(e_i)^2 + \eta(e_j)^2\} - \alpha(e_i, e_i) - \alpha(e_j, e_j) \\ &= R(e_i, e_j, e_j, e_i) + g(h(e_i, e_j), h(e_i, e_j)) - g(h(e_i, e_i), h(e_j, e_j)). \end{aligned}$$

By summation after  $1 \leq i, j \leq n$ , it follows from the previous relation that

$$(3.7) \quad \begin{aligned} 2\tau + \|h\|^2 - n^2 \|H\|^2 &= -2(n-1)\lambda + (n^2 - n)\frac{c}{4} \\ &\quad + \frac{3c}{4} \|P\|^2 - \frac{c}{2}(n-1)\|\xi^\top\|^2. \end{aligned}$$

We take

$$(3.8) \quad \begin{aligned} \varepsilon &= 2\tau - \frac{n^2(n-2)}{n-1} \|H\|^2 + 2(n-1)\lambda - (n^2 - n)\frac{c}{4} \\ &\quad - \frac{3c}{4} \|P\|^2 + \frac{c}{2}(n-1)\|\xi^\top\|^2. \end{aligned}$$

Then, from (3.7) and (3.8) we get

$$(3.9) \quad n^2 \|H\|^2 = (n-1) \left( \|h\|^2 + \varepsilon \right).$$

Let  $x \in M^n, \pi \subset T_x M^n, \dim \pi = 2, \pi = \text{span}\{e_1, e_2\}$ . We define  $e_{n+1} = \frac{H}{\|H\|}$  and from the relation (3.9) we obtain:

$$\left( \sum_{i=1}^n h_{ii}^{n+1} \right)^2 = (n-1) \left( \sum_{i,j=1}^n \sum_{r=n+1}^{2m+1} (h_{ij}^r)^2 + \varepsilon \right),$$

or equivalently,

$$\left(\sum_{i=1}^n h_{ii}^{n+1}\right)^2 = (n-1) \left[ \sum_{i=1}^n (h_{ii}^{n+1})^2 + \sum_{i \neq j} (h_{ij}^{n+1})^2 + \sum_{i,j=1}^n \sum_{r=n+2}^{2m+1} (h_{ij}^r)^2 + \varepsilon \right].$$

By using the algebraic Lemma we have from the previous relation

$$2h_{11}^{n+1}h_{22}^{n+1} \geq \sum_{i \neq j} (h_{ij}^{n+1})^2 + \sum_{i,j=1}^n \sum_{r=n+2}^{2m+1} (h_{ij}^r)^2 + \varepsilon.$$

If we denote by  $\xi_\pi = pr_\pi \xi$  we can write (see [19])

$$-\eta(e_1)^2 - \eta(e_2)^2 = -\|\xi_\pi\|^2.$$

The Gauss equation for  $X = W = e_1, Y = Z = e_2$  gives

$$\begin{aligned} K(\pi) &= R(e_1, e_2, e_2, e_1) = \frac{c}{4} + \frac{3c}{4}g^2(Pe_1, e_2) - \frac{c}{4}\|\xi_\pi\|^2 \\ &\quad - \alpha(e_1, e_1) - \alpha(e_2, e_2) + \sum_{r=n+1}^{2m+1} [h_{11}^r h_{22}^r - (h_{12}^r)^2] \\ &\geq \frac{c}{4} + \frac{3c}{4}g^2(Pe_1, e_2) - \frac{c}{4}\|\xi_\pi\|^2 - \alpha(e_1, e_1) - \alpha(e_2, e_2) \\ &\quad + \frac{1}{2} \left[ \sum_{i \neq j} (h_{ij}^{n+1})^2 + \sum_{i,j=1}^n \sum_{r=n+2}^{2m+1} (h_{ij}^r)^2 + \varepsilon \right] \\ &\quad + \sum_{r=n+2}^{2m+1} h_{11}^r h_{22}^r - \sum_{r=n+1}^{2m+1} (h_{12}^r)^2 \\ &= \frac{c}{4} + \frac{3c}{4}g^2(Pe_1, e_2) - \frac{c}{4}\|\xi_\pi\|^2 - \alpha(e_1, e_1) - \alpha(e_2, e_2) \\ &\quad + \frac{1}{2} \sum_{i \neq j} (h_{ij}^{n+1})^2 + \frac{1}{2} \sum_{i,j=1}^n \sum_{r=n+2}^{2m+1} (h_{ij}^r)^2 + \frac{1}{2}\varepsilon \\ &\quad + \sum_{r=n+2}^{2m+1} h_{11}^r h_{22}^r - \sum_{r=n+1}^{2m+1} (h_{12}^r)^2 \\ &= \frac{c}{4} + \frac{3c}{4}g^2(Pe_1, e_2) - \frac{c}{4}\|\xi_\pi\|^2 - \alpha(e_1, e_1) - \alpha(e_2, e_2) \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2} \sum_{i \neq j} (h_{ij}^{n+1})^2 + \frac{1}{2} \sum_{r=n+2}^{2m+1} \sum_{i,j>2} (h_{ij}^r)^2 + \frac{1}{2} \sum_{r=n+2}^{2m+1} (h_{11}^r + h_{22}^r)^2 \\
& + \sum_{j>2} [(h_{1j}^{n+1})^2 + (h_{2j}^{n+1})^2] + \frac{1}{2} \varepsilon \\
& \geq \frac{c}{4} + \frac{3c}{4} g^2(Pe_1, e_2) - \frac{c}{4} \|\xi_\pi\|^2 - \alpha(e_1, e_1) - \alpha(e_2, e_2) + \frac{\varepsilon}{2},
\end{aligned}$$

which implies

$$K(\pi) \geq \frac{c}{4} + \frac{3c}{4} g^2(Pe_1, e_2) - \frac{c}{4} \|\xi_\pi\|^2 - \alpha(e_1, e_1) - \alpha(e_2, e_2) + \frac{\varepsilon}{2}.$$

Denote by

$$\alpha(e_1, e_1) + \alpha(e_2, e_2) = \lambda - \text{trace}(\alpha_{|\pi^\perp}),$$

(see [19]). From (3.8) it follows

$$\begin{aligned}
K(\pi) & \geq \tau - (n-2) \left[ \frac{n^2}{2(n-1)} \|H\|^2 + (n+1) \frac{c}{8} - \lambda \right] \\
& + \frac{c}{4} \left( 3\Theta^2(\pi) - \frac{3}{2} \|P\|^2 + (n-1) \|\xi^\top\|^2 - \|\xi_\pi\|^2 \right) + \text{trace}(\alpha_{|\pi^\perp}),
\end{aligned}$$

which represents the inequality to prove.  $\square$

**Corollary 3.2.** *Under the same assumptions as in Theorem 3.1 if  $\xi$  is tangent to  $M^n$ , we have*

$$\begin{aligned}
\tau(x) - K(\pi) & \leq (n-2) \left[ \frac{n^2}{2(n-1)} \|H\|^2 + (n+1) \frac{c}{8} - \lambda \right] \\
& - \frac{c}{4} \left( 3\Theta^2(\pi) - \frac{3}{2} \|P\|^2 + n-1 - \|\xi_\pi\|^2 \right) - \text{trace}(\alpha_{|\pi^\perp}).
\end{aligned}$$

If  $\xi$  is normal to  $M^n$ , we have

$$\begin{aligned}
\tau(x) - K(\pi) & \leq (n-2) \left[ \frac{n^2}{2(n-1)} \|H\|^2 + (n+1) \frac{c}{8} - \lambda \right] \\
& - \frac{c}{4} \left( 3\Theta^2(\pi) - \frac{3}{2} \|P\|^2 \right) - \text{trace}(\alpha_{|\pi^\perp}).
\end{aligned}$$

Recall the following important result (Proposition 1.2) from [12].

**Proposition 3.3.** *The mean curvature  $H$  of  $M^n$  with respect to the semi-symmetric metric connection coincides with the mean curvature  $\overset{\circ}{H}$  of  $M^n$  with respect to the Levi-Civita connection if and only if the vector field  $U$  is tangent to  $M^n$ .*

**Remark 3.4.** According to the formula (7) from [20] (see also Proposition 3.3), it follows that  $h = \overset{\circ}{h}$  if  $U$  is tangent to  $M^n$ . In this case inequality (3.4) becomes

$$\begin{aligned} \tau(x) - K(\pi) \leq (n-2) \left[ \frac{n^2}{2(n-1)} \left\| \overset{\circ}{H} \right\|^2 + (n+1) \frac{c}{8} - \lambda \right] \\ - \frac{c}{4} \left( 3\Theta^2(\pi) - \frac{3}{2} \|P\|^2 + (n-1) \|\xi^\top\|^2 - \|\xi_\pi\|^2 \right) - \text{trace}(\alpha_{|\pi^\perp}). \end{aligned}$$

**Proposition 3.5.** *If the vector field  $U$  is tangent to  $M^n$ , then the equality case of inequality (3.4) holds at a point  $x \in M^n$  if and only if there exists an orthonormal basis  $\{e_1, e_2, \dots, e_n\}$  of  $T_x M^n$  and an orthonormal basis  $\{e_{n+1}, \dots, e_{n+p}\}$  of  $T_x^\perp M^n$  such that the shape operators of  $M^n$  in  $N^{2m+1}(c)$  at  $x$  have the following forms:*

$$A_{e_{n+1}} = \begin{pmatrix} a & 0 & 0 & \cdots & 0 \\ 0 & b & 0 & \cdots & 0 \\ 0 & 0 & \mu & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \mu \end{pmatrix}, \quad a + b = \mu,$$

$$A_{e_r} = \begin{pmatrix} h_{11}^r & h_{12}^r & 0 & \cdots & 0 \\ h_{12}^r & -h_{11}^r & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix}, \quad n+2 \leq r \leq 2m+1,$$

where we denote by  $h_{ij}^r = g(h(e_i, e_j), e_r)$ ,  $1 \leq i, j \leq n$  and  $n+2 \leq r \leq 2m+1$ .

**Proof.** The equality case holds at a point  $x \in M^n$  if and only if it achieves the equality in all the previous inequalities and we have the equality

in the Lemma.

$$\begin{aligned} h_{ij}^{n+1} &= 0, \forall i \neq j, i, j > 2, \\ h_{ij}^r &= 0, \forall i \neq j, i, j > 2, r = n + 1, \dots, 2m + 1, \\ h_{11}^r + h_{22}^r &= 0, \forall r = n + 2, \dots, 2m + 1, \\ h_{1j}^{n+1} &= h_{2j}^{n+1} = 0, \forall j > 2, \\ h_{11}^{n+1} + h_{22}^{n+1} &= h_{33}^{n+1} = \dots = h_{nn}^{n+1}. \end{aligned}$$

We may chose  $\{e_1, e_2\}$  such that  $h_{12}^{n+1} = 0$  and we denote by  $a = h_{11}^r$ ,  $b = h_{22}^r$ ,  $\mu = h_{33}^{n+1} = \dots = h_{nn}^{n+1}$ .  $\square$

It follows that the shape operators take the desired forms.

#### 4. $k$ -Ricci curvature for submanifolds of cosymplectic space forms

We first state a relationship between the sectional curvature of a submanifold  $M^n$  of a cosymplectic space form  $N^{2m+1}(c)$  of constant  $\varphi$ -sectional curvature  $c$  endowed with a semi-symmetric metric connection  $\tilde{\nabla}$  and the squared mean curvature  $\|H\|^2$ . Using this inequality, we prove a relationship between the  $k$ -Ricci curvature of  $M^n$  (intrinsic invariant) and the squared mean curvature  $\|H\|^2$  (extrinsic invariant), as another answer of the basic problem in submanifold theory which we have mentioned in the introduction.

In this section we suppose that the vector field  $U$  is tangent to  $M^n$ .

**Theorem 4.1.** *Let  $M^n, n \geq 3$ , be an  $n$ -dimensional submanifold of an  $(2m + 1)$ -dimensional a cosymplectic space form  $N^{2m+1}(c)$  of constant  $\varphi$ -sectional curvature  $c$  endowed with a semi-symmetric metric connection  $\tilde{\nabla}$  such that the vector field  $U$  is tangent to  $M^n$ . Then we have*

$$(4.1) \quad \|H\|^2 \geq \frac{2\tau}{n(n-1)} + \frac{2}{n}\lambda - \frac{c}{4} - \frac{3c}{4n(n-1)} \|P\|^2 + \frac{c}{2n} \|\xi^\top\|^2.$$

**Proof.** Let  $x \in M^n$  and  $\{e_1, e_2, \dots, e_n\}$  and orthonormal basis of  $T_x M^n$ . The relation (3.7) is equivalent with

$$(4.2) \quad n^2 \|H\|^2 = 2\tau + \|h\|^2 + 2(n-1)\lambda - (n^2 - n)\frac{c}{4} - \frac{3c}{4} \|P\|^2 + \frac{c}{2}(n-1)\|\xi^\top\|^2.$$

We choose an orthonormal basis  $\{e_1, \dots, e_n, e_{n+1}, \dots, e_{n+p}\}$  at  $x$  such that  $e_{n+1}$  is parallel to the mean curvature vector  $H(x)$  and  $e_1, \dots, e_n$  diagonalize the shape operator  $A_{e_{n+1}}$ . Then the shape operators take the forms

$$A_{e_{n+1}} \begin{pmatrix} a_1 & 0 & \dots & 0 \\ 0 & a_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & a_n \end{pmatrix},$$

$$A_{e_r} = (h_{ij}^r), \quad i, j = 1, \dots, n; \quad r = n + 2, \dots, 2m + 1, \quad \text{trace } A_{e_r} = 0.$$

From (4.2), we get

$$(4.3) \quad \begin{aligned} n^2 \|H\|^2 &= 2\tau + \sum_{i=1}^n a_i^2 + \sum_{r=n+2}^{2m+1} \sum_{i,j=1}^n (h_{ij}^r)^2 + 2(n-1)\lambda \\ &\quad - (n^2 - n)\frac{c}{4} - \frac{3c}{4} \|P\|^2 + \frac{c}{2}(n-1)\|\xi^\top\|^2. \end{aligned}$$

Since  $\sum_{i=1}^n a_i^2 \geq n \|H\|^2$ , hence we obtain

$$n^2 \|H\|^2 \geq 2\tau + n \|H\|^2 + 2(n-1)\lambda - (n^2 - n)\frac{c}{4} - \frac{3c}{4} \|P\|^2 + \frac{c}{2}(n-1)\|\xi^\top\|^2.$$

Last inequality represents (4.1). □

Using Theorem 4.1, we obtain the following result:

**Theorem 4.2.** *Let  $M^n, n \geq 3$ , be an  $n$ -dimensional submanifold of an  $(2m + 1)$ -dimensional cosymplectic space form  $N^{2m+1}(c)$  of constant  $\varphi$ -sectional curvature  $c$  endowed with a semi-symmetric metric connection  $\bar{\nabla}$ , such that the vector field  $U$  is tangent to  $M^n$ . Then, for any integer  $k, 2 \leq k \leq n$ , and any point  $x \in M^n$ , we have*

$$(4.4) \quad \|H\|^2(x) \geq \Theta_k(x) + \frac{2}{n}\lambda - \frac{c}{4} - \frac{3c}{4n(n-1)} \|P\|^2 + \frac{c}{2n}\|\xi^\top\|^2.$$

**Proof.** Let  $\{e_1, \dots, e_n\}$  be an orthonormal basis of  $T_x M$ . Denote by  $L_{i_1 \dots i_k}$  the  $k$ -plane section spanned by  $e_{i_1}, \dots, e_{i_k}$ . By the definitions, one has

$$(4.5) \quad \tau(L_{i_1 \dots i_k}) = \frac{1}{2} \sum_{i \in \{i_1, \dots, i_k\}} Ric_{L_{i_1 \dots i_k}}(e_i),$$

$$(4.6) \quad \tau(x) = \frac{1}{C_{n-2}^{k-2}} \sum_{1 \leq i_1 < \dots < i_k \leq n} \tau(L_{i_1 \dots i_k}).$$

From (4.1), (4.5) and (4.6), one derives  $\tau(x) \geq \frac{n(n-1)}{2}\Theta_k(x)$ , which implies (4.4).  $\square$

## REFERENCES

1. ALEGRE, P.; CARRIAZO, A.; KIM, Y.H.; YOON, D.W. – *B.Y. Chen's inequality for submanifolds of generalized space forms*, Indian J. Pure Appl. Math., 38 (2007), 185–201.
2. BLAIR, D.E. – *Riemannian Geometry of Contact and Symplectic Manifolds*, Progress in Mathematics, 203, Birkhäuser Boston, Inc., Boston, MA, 2002.
3. ARSLAN, K.; EZENTAS, R.; MIHAI, I.; MURATHAN, C.; ÖZGÜR, C. – *B. Y. Chen inequalities for submanifolds in locally conformal almost cosymplectic manifolds*, Bull. Inst. Math. Acad. Sinica, 29 (2001), 231–242.
4. ARSLAN, K.; EZENTAS, R.; MIHAI, I.; MURATHAN, C.; ÖZGÜR, C. – *Certain inequalities for submanifolds in  $(k, \mu)$ -contact space forms*, Bull. Austral. Math. Soc., 64 (2001), 201–212.
5. CHEN, B.-Y. – *Some pinching and classification theorems for minimal submanifolds*, Arch. Math. (Basel), 60 (1993), 568–578.
6. CHEN, B.-Y. – *Strings of Riemannian invariants, inequalities, ideal immersions and their applications*, The Third Pacific Rim Geometry Conference (Seoul, 1996), 7–60, Monogr. Geom. Topology, 25, Int. Press, Cambridge, MA, 1998.
7. CHEN, B.-Y. – *Relations between Ricci curvature and shape operator for submanifolds with arbitrary codimensions*, Glasg. Math. J., 41 (1999), 33–41.
8. CHEN, B.-Y. –  *$\delta$ -Invariants, Inequalities of Submanifolds and Their Applications*, Topics in differential geometry, 29–155, Ed. Acad. Române, Bucharest, 2008.
9. CHEN, B.-Y. – *Pseudo-Riemannian Geometry,  $\delta$ -Invariants and Applications*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2011.
10. FRIEDMANN, A.; SCHOUTEN, J.A. – *Über die Geometrie der halbsymmetrischen Übertragungen*, Math. Z., 21 (1924), 211–223.
11. HAYDEN, H.A. – *Subspaces of a space with torsion*, Proc. London Math. Soc., 34 (1932), 27–50.
12. IMAI, T. – *Hypersurfaces of a Riemannian manifold with semi-symmetric metric connection*, Tensor (N.S.), 23 (1972), 300–306.
13. IMAI, T. – *Notes on semi-symmetric metric connections*, Vol. I. Tensor (N.S.), 24 (1972), 293–296.
14. KIM, J.-S.; CHOI, J. – *A basic inequality for submanifolds in a cosymplectic space form*, Int. J. Math. Math. Sci., 9 (2003), 539–547.

15. LUDDEN, G.D. – *Submanifolds of cosymplectic manifolds*, J. Differential Geometry, 4 (1970), 237–244.
16. MATSUMOTO, K.; MIHAI, I.; OIAGĂ, A. – *Ricci curvature of submanifolds in complex space forms*, Rev. Roumaine Math. Pures Appl., 46 (2001), 775–782.
17. MIHAI, A. – *Modern Topics in Submanifold Theory*, Editura Universității București, Bucharest, 2006.
18. MIHAI, A.; ÖZGÜR, C. – *Chen inequalities for submanifolds of real space forms with a semi-symmetric metric connection*, Taiwanese J. Math., 14 (2010), 1465–1477.
19. MIHAI, A.; ÖZGÜR, C. – *Chen inequalities for submanifolds of complex space forms and Sasakian space forms endowed with semi-symmetric metric connections*, Rocky Mountain J. Math., 41 (2011), 1653–1673.
20. NAKAO, Z. – *Submanifolds of a Riemannian manifold with semisymmetric metric connections*, Proc. Amer. Math. Soc., 54 (1976), 261–266.
21. OIAGA, A.; MIHAI, I. – *B. Y. Chen inequalities for slant submanifolds in complex space forms*, Demonstratio Math., 32 (1999), 835–846.
22. YANO, K. – *On semi-symmetric metric connection*, Rev. Roumaine Math. Pures Appl., 15 (1970), 1579–1586.
23. YOON, D.W. – *Inequality for Ricci curvature of slant submanifolds in cosymplectic space forms*, Turkish J. Math., 30 (2006), 43–56.

Received: 26.I.2010

Department of Mathematics,  
Balıkesir University,  
10145, Çağış, Balıkesir, Turkey  
TURKEY  
cozgur@@balikesir.edu.tr

Uludağ University,  
Department of Mathematics,  
16059, Görükle, Bursa,  
TURKEY  
cengiz@@uludag.edu.tr